

### Menu Objective 6

Use certified EHR technology to identify patient-specific education resources and provide to patient - **170.302(m)**

More than 10% of all unique patients seen by the EP are provided patient-specific education resources

eClinicalWorks gives providers the option to create, review, update, or delete patient education materials. Practices can choose from three (3) vendors of patient education materials – A.D.A.M., Krames Exit-Writer, or Healthwise. The practice can also create their own patient education materials and save them to a local file. The Treatment screen opens a browser mapped to the specific location for easy retrieval of the documentation (Custom Patient Education button). Medication information is provided by the integrated Medi-Span Drug Reference database.

**Rx Education**

Read this medicine information sheet carefully each time you get this medicine filled.  
This medicine information sheet has been written specifically for female patients.

**Budesonide/Formoterol Inhalation Aerosol**  
**Pronunciation:** bue-DES-oh-nide/for-MOE-ter-ol  
**Brand Name:** Symbicort  
**Product Dispensed:** Symbicort

Long-acting beta-agonists such as one of the active ingredients in this medicine (formoterol) have been rarely associated with an increased risk of asthma-related death. They may also increase the risk of asthma-related hospitalization in CHILDREN and TEENAGERS. This medicine should only be used to treat asthma patients who are not controlled by using another long-term asthma-control medicine. This medicine should not be used in patients whose asthma is well controlled by inhaled corticosteroids.

**This medicine is used for:**

Treating asthma in certain patients. It may be used for long-term treatment of chronic obstructive pulmonary disease (COPD). It may also be used for other conditions as determined by your doctor.

This medicine is a corticosteroid and a long-acting beta-agonist bronchodilator combination. It works by reducing inflammation and widening the airways in the lungs, which helps you breathe more easily.

**Do NOT use this medicine if:**

- you are allergic to any ingredient in this medicine
- you are using another medicine that has a long-acting beta-agonist (eg, salmeterol) in it
- you are having an asthma attack or severe breathing problems (eg, sudden, severe onset or worsening of asthma or COPD symptoms, such as wheezing, cough, chest tightness, shortness of breath)
- you are taking mifepristone
- you have asthma that is already well controlled with the use of a long-term asthma-control medicine (eg, inhaled corticosteroid)

Contact your doctor or health care provider right away if any of these apply to you.

**Before using this medicine:**

Print Preview... Print Close